ELSEVIER

Journal of Organometallic Chemistry 541 (1997) 473–474

Journal of Organometallic Chemistry

Author Index of Volume 541

Airola, K., 237
Akhnoukh, T., 207
Aldinger, F., 345
Allmann, C., 291
Alt, H.G., 3
Aumann, R., 187

Baratta, W., 445
Barz, M., 39
Beck, W., 423
Berke, H., 19
Bianchini, C., 143
Bill, J., 345
Blüchel, C., 89
Bock, M., 465
Boersma, J., 181
Borgmann, C., 367
Braun, T., 127
Braunstein, P., 417
Brintzinger, H.H., 219
Brunner, H., 89
Buchmeiser, M., 249
Butchard, J.R., 407

Carty, A.J., 461
Casares, J.A., 143
Cauzzi, D., 377
Costa, M., 377
Curnow, O.J., 407

Dahlenburg, L., 465
Daniel, T., 127
Dauer, B., 355
Dinjus, E., 243
Dorer, B., 219
Dötz, K.H., 285
Doyle, M.P., 89
Driediger, J., 461
Driess, A., 367
Dürr, U., 307

Eichbichler, J., 233
Enright, G.D., 461
Escarpa Gaede, P., 207

Fässler, T.F., 269
Fawzi, R., 167
Fehlhammer, W.P., 51
Fehlner, T.P., 57
Fenske, D., 423
Filippou, A.C., 333
Fischer, H., 321
Fröhlich, R., 187

Gauss, C., 19
Geiß, A., 441
Gevert, O., 127
Gomes, P.T., 121
Gonsalvi, L., 377
Graiff, C., 417
Green, M.L.H., 121
Guo, A.-I., 207

Haase, W., 57
Hahn, F.E., 51
Haning, H., 117
Hartmann, M., 261
Haug, R., 345
Hautot, D., 57
Hecht, J., 187
Heck, J., 321
Heckmann, G., 261
Heinemann, F., 291
Heinemann, F.W., 307
Herberhold, M., 391
Herdtweck, E., 445
Herminghaus, S., 321
Herrmann, W.A., 445
Hinz, I., 333
Hofmann, J., 199
Hohmann, F., 285
Holzhey, N., 243
Houbrechts, S., 321
Huttenloch, M.E., 219

Imhof, W., 109

Jaitner, P., 71, 249
Jany, G., 363
Jaouen, G., 355
Jastrzebski, J.T.B.H., 181
Jutzi, P., 9

Keller, M., 441
Kernbach, U., 51
Kisch, H., 307
Klewitz, S., 321
Knoch, F., 291
Kociok-Köhn, G., 333
Kooijman, H., 181
Köster, R., 97
Kreißl, F.R., 71
Kubiak, C.P., 277
Kunkely, H., 177

Lang, H., 157
Le Floch, P., 277
Lehotkay, Th., 71
Lei, X., 57
Leibfritz, T., 167
Leskelä, M., 363
Limberg, C., 367
Lindner, E., 167
Long, G.J., 57
Lügger, T., 51

Maisel, H.E., 97
Malleier, R., 233
Martins, A.M., 121
Masi, D., 143
Mathey, F., 277
Meli, A., 143
Meyer, T., 321
Mézailles, N., 277
Mieling, I., 9
Mihan, S., 423
Milius, W., 3, 97
Moran, P., 207
Morise, X., 417
Mountford, P., 121
Müller, J., 207
Müller, K., 345

Neumann, B., 9
Niecke, E., 237
Nieger, M., 237, 285
Nuber, B., 157

Ogric, C., 71
Orama, O., 19

Pellinghelli, M.A., 377
Peringer, P., 233
Pietschnig, R., 237
Pitter, S., 243
Podschadly, O., 321
Pohl, W., 143
Polamo, M., 363
Predieri, G., 377
Prosenc, M.-H., 219

Qiao, K., 207

Elsevier Science S.A.

Reetz, M.T., 117
Repo, T., 363
Ricard, L., 277
Rief, U., 219
Rieker, C., 249
Rijnberg, E., 181
Rosoff, M., 77
Roth, G., 321
Roths, K., 187
Rudler, H., 77
Rudler, M., 77

Salo, M., 363
Schmid, M.A., 3
Schmidt, K., 219
Schottenberger, H., 249
Schubert, S., 199
Schuhmacher, J., 345
Schulz, M., 127
Schütz, U., 269
Schwarz, W., 261
Schwarze, B., 97
Seeger, S., 401
Sellmann, D., 291
Shang, M., 57
Siemoneit, S., 285
Smail, S.J., 407
Spek, A.L., 181
Spiegler, M., 39
Stammler, H.-G., 9
Steimann, M., 167
Sünkel, K., 199
Sutter, J., 291

Thiel, W.R., 39
Thurner, C.L., 39
Tiripicchio, A., 377, 417
Top, S., 355
Tröbs, V., 391

Ullrich, B., 401

Vahrenkamp, H., 441
Vaissermann, J., 77, 355
Van Koten, G., 181
Veghini, D., 19
Vizza, F., 143
Vogler, A., 177

Wang, W., 461
Waschbüsch, K., 277
Weber, C., 157
Weidmann, T., 423
Weinmann, M., 345
Weinmann, S., 157
Weinrich, V., 423
Weiß, A., 465
Werner, H., 127
Werner, R., 57
Westerhausen, M., 261
Wolfrum, J., 401
Wössner, D., 333
Wrackmeyer, B., 97, 391
Wu, I.-Y., 157
Wurst, K., 71, 233, 249

Yu, Z., 187

Zahl, A., 465
Zanoni, R., 377

ELSEVIER

Journal of Organometallic Chemistry 541 (1997) 475–482

Journal of Organometallic Chemistry

Subject Index of Volume 541

Absorption spectra
Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177
Alkoxysilane
Komplexchemie perhalogenierter Cyclopentadiene und Alkine, XIX. Untersuchungen zur Darstellung und Reaktivität von Cymantrenyl-alkoxysilanen $[C_5X_{5-n}(SiMe_2OR)_n]Mn(CO)_3$ ($n = 1, 2$; X = Cl, Br, H) (K. Sünkel, S. Schubert und J. Hofmann), 199
Alkyne complexes
Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285
Alkynediide
Kohlenwasserstoffverbrückte Metallkomplexe, XXXIX. Zur Chemie von Pentacarbonylrheniumacetyliden (S. Mihan, T. Weidmann, V. Weinrich, D. Fenske and W. Beck), 423
Alkyne insertion
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
Alkynes
π-Olefin–Iridium-Komplexe XXIV. Photoreaktionen von $[CpM(C_2H_4)_2]$ (M = Rh, Ir) mit Alkinen — Fundgrube für neue π-Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-l. Guo, P. Moran und K. Qiao), 207
Alkynyl ligand
Kohlenwasserstoffverbrückte Metallkomplexe, XXXIX. Zur Chemie von Pentacarbonylrheniumacetyliden (S. Mihan, T. Weidmann, V. Weinrich, D. Fenske and W. Beck), 423
Allylic rearrangement
1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
Allylindium reagents
Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117
Aminocarbene complexes
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187
Anionic complex
Synthesis and structural characterization of the homoleptic magnesium bis zincate complex $Mg(thf)_6[Zn(CH_2Ph)_3]_2$ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181
Anti-tumour
Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann and G. Jaouen), 355
Aryloxides
Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363

Bisoxazoline
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
Bis[(σ)-1,3-butadiyne-1-yl] titanocenes
Synthese und Reaktionsverhalten der Titanocen-Verbindungen $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CR)_2$ und $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}SiMe_2{-}C{\equiv}CSiMe_3)_2$; Kristallstruktur von $[(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CC_2H_5)_2]Pd(PPh_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
Bis[(σ)-1,4-pentadiyne-1-yl] titanocenes
Synthese und Reaktionsverhalten der Titanocen-Verbindungen $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CR)_2$ und $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}SiMe_2{-}C{\equiv}CSiMe_3)_2$; Kristallstruktur von $[(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CC_2H_5)_2]Pd(PPh_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
1,3-Butadienes
1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
Butadiynyl
Ethinyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $Cp^*Fe(Ph_2PCH(X)CH_2PPh_2)C{\equiv}CY$ (X = H, PPh_2; Y = H, PPh_2, $P^{(+)}Ph_2Me$) und $Ru(Ph_2PCH_2PPh_2)_2(X)C{\equiv}CC{\equiv}CSiMe_3$ (X = Cl, $C{\equiv}CC{\equiv}CSiMe_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Calcium
Synthese und Struktur von Bis[η^5-(cyclopropyl-cyclopropylidenmethyl)cylopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. Westerhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
Calcocen
Synthese und Struktur von Bis[η^5-(cyclopropyl-cyclopropylidenmethyl)cylopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. Westerhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
Carbene complexes
Metal–carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187

Elsevier Science S.A.

Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diiodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51
Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285

Carbene ligand
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente [M('S_2C')] [M = Ni^{II}, Pd^{II}, Pt^{II}; '$S_2C'^{2-}$ = 1,3-Imidazolidinyl-*N*,*N'*-bis(2-benzolthiolat)(2 –)]. [M(L)('S_2C')]-Komplexe mit L = CN^-, CH_3^-, $COCH_3^-$, CNBu, CR_2, SH^- und SR^- (R = Et, Ph, *o*-$C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291

Carbene ligands
Cyclopentadienyl– und Pentamethylcyclopentadienyl–Rutheniumkomplexe mit Carboxylat-, Vinylester- und Carben-Liganden (H. Werner, T. Braun, T. Daniel, O. Gevert und M. Schulz), 127

Carbon dioxide
Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO_2 mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243

Carbonyls
Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277

Carboxylate
Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron–cobalt and manganese–cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57

Carboxylathoruthenium(II) complexes
Cyclopentadienyl– und Pentamethylcyclopentadienyl–Rutheniumkomplexe mit Carboxylat-, Vinylester- und Carben-Liganden (H. Werner, T. Braun, T. Daniel, O. Gevert und M. Schulz), 127

Carbyne complexes
Metal–carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333

C–H activation
Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-(Fc)N–CH}_2\text{–}\overline{C = C\text{–}C(H) = C(H)\text{–}N}\text{Me}]\text{-}Fe_2(CO)_6$ (Fc = $(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$) (W. Imhof), 109

Chelate complexes
Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285

Chelate ligands
Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39

Chemoselectivity
Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117

Chromium
Metal–carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
Synthesis and NMR spectroscopy of [t-$Bu_2PHg(\mu\text{-}PPh_2)M(CO)_5$], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233

Chromium and tungsten complexes
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187

Chromium complexes
Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285

Cluster
Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[Co_3(CO)_9\{\eta^1\text{-}\mu_3\text{-}[CP(O)(OEt)_2]\}]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-(Fc)N–CH}_2\text{–}\overline{C = C\text{–}C(H) = C(H)\text{–}N}\text{Me}]\text{-}Fe_2(CO)_6$ (Fc = $(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$) (W. Imhof), 109

Cobalt
Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[Co_3(CO)_9\{\eta^1\text{-}\mu_3\text{-}[CP(O)(OEt)_2]\}]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
Synthese und Reaktionsverhalten der Titanocen-Verbindungen $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C\text{–}C{\equiv}CR)_2$ und $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C\text{–}SiMe_2\text{–}C{\equiv}CSiMe_3)_2$; Kristallstruktur von $[(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C\text{–}C{\equiv}CC_2H_5)_2]Pd(PPh_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157

Cobalt catalysis
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307

Cobalt cluster
Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron–cobalt and manganese–cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57

Conjugated π systems
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187

Co-oligomerisation
Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO_2 mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243

Copper
Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277

Copper catalysts
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
Coupling constant sign determination
Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
Crystal structure
Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377
Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron–cobalt and manganese–cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57
Synthesis and X-ray structures of the five-coordinate zirconocene complexes $PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2ZrCl_2$, $[PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2ZrCl(H_2O)]Cl$ and $PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2Zr(NCS)_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407
Crystal structures
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente $[M(\text{'}S_2C\text{'})]$ $[M = Ni^{II}, Pd^{II}, Pt^{II}; \text{'}S_2C\text{'}^{2-} = 1,3\text{-Imidazolidinyl-}N,N'\text{-bis(2-benzolthiolat)}(2-)]$. $[M(L)(\text{'}S_2C\text{'})]$-Komplexe mit $L = CN^-, CH_3^-, COCH_3^-$, CNBu, CR_2, SH^- und SR^- (R = Et, Ph, $o\text{-}C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
C–S activation
Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
Cyclic voltammetry
Metal–carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333
Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
Cycloaddition
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
Cyclobutenylidene complexes
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
Cyclopentadienyl
Komplexchemie perhalogenierter Cyclopentadiene und Alkine, XIX. Untersuchungen zur Darstellung und Reaktivität von Cymantrenyl-alkoxysilanen $[C_5X_{5-n}(SiMe_2OR)_n]Mn(CO)_3$ (n = 1, 2; X = Cl, Br, H) (K. Sünkel, S. Schubert und J. Hofmann), 199

Diaryldiazene
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
Di(cyclopropyl)fulvene
Synthese und Struktur von Bis[η^5-(cyclopropyl-cyclopropylidenmethyl)cylopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. Westerhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
Diesters
Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167
2,3-Dihydrocinnoline
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
Double-bridged cyclopentadienyl
$\pi\text{-}H_3C_5(Me_2Si)(Me_2SiSiMe_2)\text{-}\pi\text{-}C_5H_3$: Ein neuartiger, zweifach verbrückter Dicyclopentadienyl-Ligand: Synthese und beispielhafter Einsatz in der Komplexchemie (P. Jutzi, I. Mieling, B. Neumann und H.-G. Stammler), 9
Dynamic molecules
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ-gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3

Enantioselectivity
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
Enantioseparation
ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219
Ethynyl
Ethinyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $Cp^*Fe(Ph_2PCH(X)CH_2PPh_2)C \equiv CY$ (X = H, PPh_2; Y = H, PPh_2, $P^{(+)}Ph_2Me$) und $Ru(Ph_2PCH_2PPh_2)_2(X)C \equiv CC \equiv CSiMe_3$ (X = Cl, C ≡ CC ≡ $CSiMe_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Ferrocene
Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann and G. Jaouen), 355
Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-}(Fc)N\text{–}CH_2\text{–}\overbrace{C = C\text{–}C(H) = C(H)}\text{–}NMe]\text{-}Fe_2(CO)_6$ ($Fc = (\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$) (W. Imhof), 109
Ferrocenylbismethylenephosphorane
Ferrocenylbis(ylene)phosphoranes (R. Pietschnig, M. Nieger, E. Niecke and K. Airola), 237
Ferrocifen
Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann and G. Jaouen), 355
η^1-Fluorenyl
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ-gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
Fluorescence immunoassay
Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Germanium
Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
Gold complexes
Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177

^{1}H NMR
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ-gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
Hydride complexes
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
Hydrodesulfurization
Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
Hydroformylation
Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377
Hydroxy
Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461

Imine
Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $[\mu_2$-η^3-(Fc)N–CH_2–$\overbrace{C=C–C(H)=C(H)}$–NMe]-$Fe_2(CO)_6$ (Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)) (W. Imhof), 109
Immobilized catalyst
Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO_2 mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243
Indoles
Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307
μ^4-Iodo
Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
Iridium
Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143
π-Olefin–Iridium-Komplexe XXIV. Photoreaktionen von [CpM(C_2H_4)$_2$] (M = Rh, Ir) mit Alkinen — Fundgrube für neue π-Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-l. Guo, P. Moran und K. Qiao), 207
Iron
Ethinyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $Cp^*Fe(Ph_2PCH(X)CH_2PPh_2)C \equiv CY$ (X = H, PPh_2; Y = H, PPh_2, $P^{(+)}Ph_2Me$) und $Ru(Ph_2PCH_2PPh_2)_2(X)C \equiv CC \equiv CSiMe_3$ (X = Cl, C ≡ CC ≡ $CSiMe_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465
Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $[\mu_2$-η^3-(Fc)N–CH_2–$\overbrace{C=C–C(H)=C(H)}$–NMe]-$Fe_2(CO)_6$ (Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)) (W. Imhof), 109
The reactions of dicarbonylbis(phosphorus donor)iron fragments with tertiary propargyl alcohols (C. Gauss, D. Veghini, O. Orama and H. Berke), 19

Ketenyl complexes
Metal–carbon multiple bonds: Half-sandwich phenylcarbyne complexes of chromium—synthesis, structure, electrochemistry and reactions with PMe_3 (A.C. Filippou, D. Wössner, G. Kociok-Köhn and I. Hinz), 333

Lead
Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391
Ligand tuning
Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117
Luminescence
Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Magnesium
Synthesis and structural characterization of the homoleptic magnesium bis zincate complex $Mg(thf)_6[Zn(CH_2Ph)_3]_2$ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181
Manganese
Komplexchemie perhalogenierter Cyclopentadiene und Alkine, XIX. Untersuchungen zur Darstellung und Reaktivität von Cymantrenyl-alkoxysilanen $[C_5X_{5-n}(SiMe_2OR)_n]Mn(CO)_3$ (n = 1, 2; X = Cl, Br, H) (K. Sünkel, S. Schubert und J. Hofmann), 199
Manganese complexes
Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167
Meerwein arylation
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89
Mercury
Synthesis and NMR spectroscopy of [t-$Bu_2PHg(\mu$-$PPh_2)M(CO)_5$], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233
Metal cubane
Clusters as ligands, part 3. Generation of tricobalt cluster carboxylate-bridged iron–cobalt and manganese–cobalt mixed-metal alkoxide cubes from iron and manganese tricobalt cluster metal carboxylates (X. Lei, M. Shang, T.P. Fehlner, R. Werner, W. Haase, D. Hautot and G.J. Long), 57
1-Metalla-1,3,5,7-octatetraenes
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187
Metallocene
ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crys-

tal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ-gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3

Metallocene complexes
π-$H_3C_5(Me_2Si)(Me_2SiSiMe_2)$-$\pi$-$C_5H_3$: Ein neuartiger, zweifach verbrückter Dicyclopentadienyl-Ligand: Synthese und beispielhafter Einsatz in der Komplexchemie (P. Jutzi, I. Mieling, B. Neumann und H.-G. Stammler), 9

Metallocenes
Synthesis and X-ray structures of the five-coordinate zirconocene complexes $PhP(CH_2CH_2$-η^5-$C_5H_4)_2ZrCl_2$, $[PhP(CH_2CH_2$-η^5-$C_5H_4)_2ZrCl(H_2O)]Cl$ and $PhP(CH_2CH_2$-η^5-$C_5H_4)_2Zr(NCS)_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407

Methyl ligand
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente [M('S_2C')] [M = Ni^{II}, Pd^{II}, Pt^{II}; '$S_2C'^{2-}$ = 1,3-Imidazolidinyl-*N*,*N*′-bis(2-benzolthiolat)(2 −)]. [M(L)('S_2C')]-Komplexe mit L = CN^-, CH_3^-, $COCH_3^-$, CNBu, CR_2, SH^- und SR^- (R = Et, Ph, *o*-$C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291

Mixed–metal bimetallocenophanes
Mixed-membered-bisfulvalene dimetal-complexes via metallocene-substituted norbornenyl alcohols (H. Schottenberger, M. Buchmeiser, C. Rieker, P. Jaitner and K. Wurst), 249

Molybdenum
Ansa-bridged η-cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121
Synthesis and NMR spectroscopy of [t-$Bu_2PHg(\mu$-$PPh_2)M(CO)_5$], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233

Molybdenum oxo compounds
Synthesis, structure and properties of bisallyltris(trimethylphosphine) molybdenum(II). X-ray crystal structures of $[Mo(\eta^1$:η^2-$C_3H_5)_2(PMe_3)_3]$ and $PMe_4^+[MoCl_4(PMe_3)_2]^-$ (C. Borgmann, C. Limberg and A. Driess), 367

Multinuclear NMR
Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391

Multiple bonds
Multiple bonds between main-group elements and transition metals: Part 157. Neutral and cationic *ansa*-metallocenes of niobium(V) and tantalum(V): synthesis, structures and stereochemical non-rigidity (W.A. Herrmann, W. Baratta and E. Herdtweck), 445

Nickel
Synthese und Reaktionsverhalten der Titanocen-Verbindungen (η^5-$C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CR)_2$ und (η^5-$C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}SiMe_2{-}C{\equiv}CSiMe_3)_2$; Kristallstruktur von [(η^5-$C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CC_2H_5)_2$]$Pd(PPh_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente [M('S_2C')] [M = Ni^{II}, Pd^{II}, Pt^{II}; '$S_2C'^{2-}$ = 1,3-Imidazolidinyl-*N*,*N*′-bis(2-benzolthiolat)(2 −)]. [M(L)('S_2C')]-Komplexe mit L = CN^-, CH_3^-, $COCH_3^-$, CNBu, CR_2, SH^- und SR^- (R = Et, Ph, *o*-$C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291

Niobium
Ansa-bridged η-cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121

Niobium (V)
Multiple bonds between main-group elements and transition metals: Part 157. Neutral and cationic *ansa*-metallocenes of niobium(V) and tantalum(V): synthesis, structures and stereochemical non-rigidity (W.A. Herrmann, W. Baratta and E. Herdtweck), 445

NLO
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

NMR
Synthese und Struktur von Bis[η^5-(cyclopropyl-cyclopropylidenmethyl)cylopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. Westerhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261

NMR spectroscopic analysis
Umsetzungen mit dem Zintl-Ion Sn_9^{4-}. Kristallstrukturanalysen von $[(\eta^5$-$C_5H_5)(CO)_2Fe]_3Sn$–OH und $K_2[K$–(2,2,2-crypt)$]_2Sn_2Te\cdot$1en. Synthese des Anions $Sn_2Te_3^{2-}$ (T.F. Fässler und U. Schütz), 269

Optically active ligands
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89

1,1-Organoboration
1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97

Organozinc
Synthesis and structural characterization of the homoleptic magnesium bis zincate complex $Mg(thf)_6[Zn(CH_2Ph)_3]_2$ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181

Palladium
Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diiodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51
Synthese und Reaktionsverhalten der Titanocen-Verbindungen (η^5-$C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CR)_2$ und (η^5-$C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}SiMe_2{-}C{\equiv}CSiMe_3)_2$; Kristallstruktur von [(η^5-$C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CC_2H_5)_2$]$Pd(PPh_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente [M('S_2C')] [M = Ni^{II}, Pd^{II}, Pt^{II}; '$S_2C'^{2-}$ = 1,3-Imidazolidinyl-*N*,*N*′-bis(2-benzolthiolat)(2 −)]. [M(L)('S_2C')]-Komplexe mit L = CN^-, CH_3^-, $COCH_3^-$, CNBu, CR_2, SH^- und SR^- (R = Et, Ph, *o*-$C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291

Pentacarbonylrhenium
Kohlenwasserstoffverbrückte Metallkomplexe, XXXIX. Zur Chemie von Pentacarbonylrheniumacetyliden (S. Mihan, T. Weidmann, V. Weinrich, D. Fenske and W. Beck), 423

Phenoxide
Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363

Phosphine
Synthesis and X-ray structures of the five-coordinate zirconocene complexes $PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2ZrCl_2$, $[PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2ZrCl(H_2O)]Cl$ and $PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2Zr(NCS)_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407
Phosphine ligands
Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39
Phosphinidene
Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461
Phosphinite ligands
Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39
2-Phosphinophosphinines
Synthesis and X-ray crystal structures of dimeric nickel(0) and tetrameric copper(I) iodide complexes of 2-diphenylphosphino-3-methylphosphinine (N. Mézailles, P. Le Floch, K. Waschbüsch, L. Ricard, F. Mathey and C.P. Kubiak), 277
Phosphonate
Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[Co_3(CO)_9\{\eta^1\text{-}\mu_3\text{-}[CP(O)(OEt)_2]\}]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
Phosphonioethynyl
Ethinyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $Cp^*Fe(Ph_2PCH(X)CH_2PPh_2)C \equiv CY$ ($X = H$, PPh_2; $Y = H$, PPh_2, $P^{(+)}Ph_2Me$) und $Ru(Ph_2PCH_2PPh_2)_2(X)C \equiv CC \equiv CSiMe_3$ ($X = Cl$, $C \equiv CC \equiv CSiMe_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465
Phosphorus
Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461
Photochemistry
π-Olefin–Iridium-Komplexe XXIV. Photoreaktionen von $[CpM(C_2H_4)_2]$ ($M = Rh$, Ir) mit Alkinen — Fundgrube für neue π-Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-l. Guo, P. Moran und K. Qiao), 207
Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177
Photoluminescence
Photochemistry of chlorocarbonyl gold(I) (H. Kunkely and A. Vogler), 177
Platinum
Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461
Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diiodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51
Synthese und Reaktionsverhalten der Titanocen-Verbindungen $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CR)_2$ und $(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}SiMe_2{-}C{\equiv}CSiMe_3)_2$; Kristallstruktur von $[(\eta^5\text{-}C_5H_4SiMe_3)_2Ti(C{\equiv}C{-}C{\equiv}CC_2H_5)_2]Pd(PPh_3)$ (H. Lang, I.-Y. Wu, S. Weinmann, C. Weber und B. Nuber), 157
Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente $[M(\text{'}S_2C\text{'})]$ [$M = Ni^{II}$, Pd^{II}, Pt^{II}; $\text{'}S_2C\text{'}^{2-}$ = 1,3-Imidazolidinyl-*N*,*N'*-bis(2-benzolthiolat)(2 −)]. $[M(L)(\text{'}S_2C\text{'})]$-Komplexe mit $L = CN^-$, CH_3^-, $COCH_3^-$, CNBu, CR_2, SH^- und SR^- ($R = Et$, Ph, $o\text{-}C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291
Polymerization
Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363
Polymer phosphines
Die heterogen katalysierte Co-oligomerisation von 1,3-Butadien und CO_2 mit immobilisierten Palladiumkomplexen (N. Holzhey, S. Pitter und E. Dinjus), 243
Polysilylcarbodi-imides
Boron-containing polysilylcarbodi-imides: a new class of molecular precursors for Si-B-C-N ceramics (M. Weinmann, R. Haug, J. Bill, F. Aldinger, J. Schuhmacher and K. Müller), 345
Pyranylidene complexes
Organic syntheses via transition metal complexes, LXXXVIII. 1-Metalla-1,3,5,7-octatetraenes from (1-alkynyl)carbene complexes of chromium and tungsten via [6-(1-alkenyl)]pyranylidene complexes (Z. Yu, R. Aumann, R. Fröhlich, K. Roths and J. Hecht), 187
Pyridinyloxazoline
Asymmetric catalysis, part 108. Copper catalysts with optically active ligands in the enantioselective Meerwein arylation of activated olefins (H. Brunner, C. Blüchel and M.P. Doyle), 89

Reductive cleavage
Mixed-membered-bisfulvalene dimetal-complexes via metallocene-substituted norbornenyl alcohols (H. Schottenberger, M. Buchmeiser, C. Rieker, P. Jaitner and K. Wurst), 249
Rhenium complexes
Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167
Rhodium
Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377
π-Olefin–Iridium-Komplexe XXIV. Photoreaktionen von $[CpM(C_2H_4)_2]$ ($M = Rh$, Ir) mit Alkinen — Fundgrube für neue π-Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-l. Guo, P. Moran und K. Qiao), 207
Ruthenium
Cyclopentadienyl– und Pentamethylcyclopentadienyl–Rutheniumkomplexe mit Carboxylat-, Vinylester- und Carben-Liganden (H. Werner, T. Braun, T. Daniel, O. Gevert und M. Schulz), 127
Ethinyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $Cp^*Fe(Ph_2PCH(X)CH_2PPh_2)C \equiv CY$ ($X = H$, PPh_2; $Y = H$, PPh_2, $P^{(+)}Ph_2Me$) und $Ru(Ph_2PCH_2PPh_2)_2(X)C \equiv CC \equiv CSiMe_3$ ($X = Cl$, $C \equiv CC \equiv CSiMe_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465
Hydroxyphosphinidene complexes. Novel organometallic cluster acids and precursors of coordinated phosphorus monoxide (W. Wang, G.D. Enright, J. Driediger and A.J. Carty), 461
Ruthenium(II) complexes
Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Saturated hydrocarbon bridges
Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167
Silica xerogels
Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the

hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377

Silicon

π-$H_3C_5(Me_2Si)(Me_2SiSiMe_2)$-$\pi$-$C_5H_3$: Ein neuartiger, zweifach verbrückter Dicyclopentadienyl-Ligand: Synthese und beispielhafter Einsatz in der Komplexchemie (P. Jutzi, I. Mieling, B. Neumann und H.-G. Stammler), 9

Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391

Silsesquioxanes

Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377

Solvatochromic effect

Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

Stepwise ring enlargement

1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97

Stereoselectivity

Ligand effects in selective addition reactions of organoindium reagents with carbonyl compounds (M.T. Reetz and H. Haning), 117

Stilbenes

Reactions of complex ligands 74. Alkyne carbene chromium chelates: Synthesis, structure and carbene dimerization (K.H. Dötz, S. Siemoneit, F. Hohmann and M. Nieger), 285

Stilbenylazobenzene

Transition metal complexes of diazenes XXXIX. Stilbenylazobenzene derivatives by cobalt-catalysed addition of diphenylacetylene to 1,2-diaryldiazenes and their acid-catalysed rearrangement to *N*-anilinoindoles (U. Dürr, F.W. Heinemann and H. Kisch), 307

Sulphur ligand

Übergangsmetallkomplexe mit Schwefelliganden CXXIV. Koordination von C- und S-Liganden an Metall–Carben–Thiolat-Fragmente [M('S_2C')] [M = Ni^{II}, Pd^{II}, Pt^{II}; '$S_2C'^{2-}$ = 1,3-Imidazolidinyl-*N*,*N*′-bis(2-benzolthiolat)(2 −)]. [M(L)('S_2C')]-Komplexe mit L = CN^-, CH_3^-, $COCH_3^-$, CNBu, CR_2, SH^- und SR^- (R = Et, Ph, *o*-$C_6H_4NH_2$) (D. Sellmann, C. Allmann, F. Heinemann, F. Knoch und J. Sutter), 291

Tamoxifen

Facile route to ferrocifen, 1-[4-(2-dimethylaminoethoxy)]-1-(phenyl-2-ferrocenyl-but-1-ene), first organometallic analogue of tamoxifen, by the McMurry reaction (S. Top, B. Dauer, J. Vaissermann and G. Jaouen), 355

Tantalum (V)

Multiple bonds between main-group elements and transition metals: Part 157. Neutral and cationic *ansa*-metallocenes of niobium(V) and tantalum(V): synthesis, structures and stereochemical non-rigidity (W.A. Herrmann, W. Baratta and E. Herdtweck), 445

Tertiary propargyl alcohols

The reactions of dicarbonylbis(phosphorus donor)iron fragments with tertiary propargyl alcohols (C. Gauss, D. Veghini, O. Orama and H. Berke), 19

Thermogravimetric analysis

Boron-containing polysilylcarbodi-imides: a new class of molecular precursors for Si-B-C-N ceramics (M. Weinmann, R. Haug, J. Bill, F. Aldinger, J. Schuhmacher and K. Müller), 345

Thiocyanate

Synthesis and X-ray structures of the five-coordinate zirconocene complexes PhP(CH_2CH_2-η^5-$C_5H_4)_2ZrCl_2$, [PhP(CH_2CH_2-η^5-$C_5H_4)_2ZrCl(H_2O)$]Cl and PhP(CH_2CH_2-η^5-$C_5H_4)_2Zr(NCS)_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407

Thiophenes

Insertion of iridium into C–H and C–S bonds of 2,5-dimethylthiophene, 2-methylbenzothiophene and 4,6-dimethyldibenzothiophene (C. Bianchini, J.A. Casares, D. Masi, A. Meli, W. Pohl and F. Vizza), 143

Thiourea ligands

Anchoring rhodium(I) on thiourea-functionalized silica xerogels and silsesquioxanes. Part II. Matrix effects on the selectivity in the hydroformylation of styrene (D. Cauzzi, M. Costa, L. Gonsalvi, M.A. Pellinghelli, G. Predieri, A. Tiripicchio and R. Zanoni), 377

Time resolved spectroscopy

Verwendung von lumineszierenden Ruthenium(II)-Komplexen zur Markierung von Antikörpern (B. Ullrich, S. Seeger und J. Wolfrum), 401

Tin

1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97

Tri(*tert*-butyl)plumbyl derivatives of Group 14 elements. Multinuclear magnetic resonance studies: determination of coupling signs (M. Herberhold, V. Tröbs and B. Wrackmeyer), 391

Titanium

Ansa-bridged η-cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121

ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219

Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of [$Co_3(CO)_9\{\eta^1$-μ_3-[CP(O)$(OEt)_2$]}] (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417

Transition metal complexes

Ligands with cycloalkane backbones. II. Chelate ligands from 2-(diphenylphosphinyl)cyclohexanol: syntheses and transition metal complexes (C.L. Thurner, M. Barz, M. Spiegler and W.R. Thiel), 39

Tungsten

Synthesis and NMR spectroscopy of [t-$Bu_2PHg(\mu$-$PPh_2)M(CO)_5$], M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233

Unsaturated hydrocarbon bridges

Dinuclear metal complexes with saturated and unsaturated hydrocarbon bridges (E. Lindner, T. Leibfritz, R. Fawzi and M. Steimann), 167

Vanadium

ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219

Vinylidene complexes

Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

X-ray analysis
1,1-Organoboration of mono-1-alkynyltin compounds using dialkyl(*N*-azolyl)boranes-stepwise ring enlargement of boron heterocycles (B. Wrackmeyer, H.E. Maisel, B. Schwarze, W. Milius and R. Köster), 97
X–ray analysis
Umsetzungen mit dem Zintl-Ion Sn_9^{4-}. Kristallstrukturanalysen von $[(\eta^5\text{-}C_5H_5)(CO)_2Fe]_3Sn\text{–}OH$ und $K_2[K\text{–}(2,2,2\text{-crypt})]_2Sn_2Te \cdot 1en$. Synthese des Anions $Sn_2Te_3^{2-}$ (T.F. Fässler und U. Schütz), 269
X-ray crystallography
Ferrocenylbis(ylene)phosphoranes (R. Pietschnig, M. Nieger, E. Niecke and K. Airola), 237
π-Olefin–Iridium-Komplexe XXIV. Photoreaktionen von $[CpM(C_2H_4)_2]$ (M = Rh, Ir) mit Alkinen — Fundgrube für neue π-Komplexe (J. Müller, T. Akhnoukh, P. Escarpa Gaede, A.-l. Guo, P. Moran und K. Qiao), 207
Platinum(II) and palladium(II) complexes with 1,2-dihydrobenzoxazol-2-ylidene ligands and molecular structure of *trans*-bis(1,2-dihydrobenzoxazol-2-ylidene)diiodo palladium (U. Kernbach, T. Lügger, F.E. Hahn and W.P. Fehlhammer), 51
Synthesis and NMR spectroscopy of $[t\text{-}Bu_2PHg(\mu\text{-}PPh_2)M(CO)_5]$, M = Cr, Mo, W and X-ray structure for M = Cr, W (J. Eichbichler, R. Malleier, K. Wurst and P. Peringer), 233
X-ray crystal structures
Synthesis, structure and properties of bisallyltris(trimethylphosphine) molybdenum(II). X-ray crystal structures of $[Mo(\eta^1\text{:}\eta^2\text{-}C_3H_5)_2(PMe_3)_3]$ and $PMe_4^+[MoCl_4(PMe_3)_2]^-$ (C. Borgmann, C. Limberg and A. Driess), 367
X-ray structural analysis
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321
X-ray structure
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ-gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
Synthese und Struktur von Bis[η^5-(cyclopropyl-cyclopropylidenmethyl)cylopentadienyl]bis(tetrahydrofuran-*O*)calcium (M. Westerhausen, M. Hartmann, G. Heckmann und W. Schwarz), 261
Synthesis and crystal structure of N-ferrocenyl substituted heterocyclic imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $[\mu_2\text{-}\eta^3\text{-}(Fc)N\text{–}CH_2\text{–}\overrightarrow{C = C\text{–}C(H) = C(H)\text{–}N}Me]\text{-}Fe_2(CO)_6$ (Fc = $(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$) (W. Imhof), 109
X-ray structure analysis
Ethinyl- und Butadiinylkomplexe des Eisens und Rutheniums mit terminalen Hauptgruppenelement-Substituenten, $Cp^*Fe(Ph_2PCH(X)CH_2PPh_2)C \equiv CY$ (X = H, PPh_2; Y = H, PPh_2, $P^{(+)}Ph_2Me$) und $Ru(Ph_2PCH_2PPh_2)_2(X)C \equiv CC \equiv CSiMe_3$ (X = Cl, C ≡ CC ≡ $CSiMe_3$) (L. Dahlenburg, A. Weiß, M. Bock und A. Zahl), 465

Ynamine
Pentacarbonyl(cyclobutenyliden)chrom-Komplexe – Synthese, Struktur und nichtlineare optische Eigenschaften (H. Fischer, O. Podschadly, G. Roth, S. Herminghaus, S. Klewitz, J. Heck, S. Houbrechts und T. Meyer), 321

Zincate
Synthesis and structural characterization of the homoleptic magnesium bis zincate complex $Mg(thf)_6[Zn(CH_2Ph)_3]_2$ (E. Rijnberg, J.T.B.H. Jastrzebski, J. Boersma, H. Kooijman, A.L. Spek and G. Van Koten), 181
Zintl anion
Umsetzungen mit dem Zintl-Ion Sn_9^{4-}. Kristallstrukturanalysen von $[(\eta^5\text{-}C_5H_5)(CO)_2Fe]_3Sn\text{–}OH$ und $K_2[K\text{–}(2,2,2\text{-crypt})]_2Sn_2Te \cdot 1en$. Synthese des Anions $Sn_2Te_3^{2-}$ (T.F. Fässler und U. Schütz), 269
Zirconium
Ansa-bridged η-cyclopentadienyl imido and amido derivatives of titanium, zirconium, niobium and molybdenum (P.T. Gomes, M.L.H. Green, A.M. Martins and P. Mountford), 121
ansa-Metallocene derivatives XXXIX. Biphenyl-bridged metallocene complexes of titanium, zirconium, and vanadium: syntheses, crystal structures and enantioseparation (M.E. Huttenloch, B. Dorer, U. Rief, M.-H. Prosenc, K. Schmidt and H.H. Brintzinger), 219
Carbonyl clusters with a capping methylidyne phosphonate ligand. Crystal structure of $[Co_3(CO)_9\{\eta^1\text{-}\mu_3\text{-}[CP(O)(OEt)_2]\}]$ (P. Braunstein, C. Graiff, X. Morise and A. Tiripicchio), 417
Sterisch überladene Zirconocenkomplexe—erstes Beispiel für einen σ-gebundenen Fluorenylliganden (M.A. Schmid, H.G. Alt und W. Milius), 3
Synthesis and X-ray structures of the five-coordinate zirconocene complexes $PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2ZrCl_2$, $[PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2ZrCl(H_2O)]Cl$ and $PhP(CH_2CH_2\text{-}\eta^5\text{-}C_5H_4)_2Zr(NCS)_2$ (J.R. Butchard, O.J. Curnow and S.J. Smail), 407
Zirconocene
Phenoxy substituted zirconocenes in ethylene polymerization (T. Repo, G. Jany, M. Salo, M. Polamo and M. Leskelä), 363